

June 3, 2016

Phil Strobel, NEPA Program Director  
Matt Hubner, NEPA Program  
Environmental Protection Agency  
NEPA Compliance and Review Program  
U.S. EPA, Region 8, 8EPR-N  
1595 Wynkoop Street  
Denver, CO 80202-1129  
Via email:  
Strobel.Philip@epa.gov  
Hubner.Matt@epa.gov

RE: Draft Environmental Impact Statement for the Enefit American Oil Utility Corridor Project.

Dear Mr. Strobel and Mr. Hubner,

Thank you for meeting with us last week to discuss the Draft Environmental Impact Statement (EIS) for the Enefit American Oil (Enefit) Utility Corridor Project, DOI-BLM-UT-G010-2014-0007-EIS (Utility Corridor DEIS). We appreciate your time, and attention to this issue.

Per our discussion, the undersigned organizations are writing to reiterate several concerns about the ongoing National Environmental Policy Act (NEPA) process for Enefit right-of-way applications. This utility corridor will enable Enefit's large scale oil shale strip mine and a 50,000 barrel per day retort operation in northeastern Utah, called the "South Project," on the Utah-Colorado border and nearby both the Green and White rivers.

If successful, Enefit's South Project will be the first commercial scale oil shale operation in the United States; thus, it is very important that the Bureau of Land Management (BLM) completes a meaningful analysis of environmental impacts – including water use, emissions, a greenhouse gas inventory, and species analysis – as part of its obligations under federal environmental laws.

We are deeply concerned that BLM has side-stepped around its obligation to take a hard look at the reasonably foreseeable impacts of the South Project in its draft environmental impact statement. We urge the Environmental Protection Agency (EPA), as a cooperating agency in the DEIS process and a primary regulator given the projects location in Indian Country, to carefully consider the following issues.

### **I. NEPA Requires Reasonable Forecasting**

In the draft EIS, BLM did not analyze the potential impacts of Enefit's South Project on the environment. Part of the stated reason for this omission is because Enefit has refused to provide a plan of development for its oil shale mining and retort operation. Enefit has withheld this information based on a claim that it cannot determine the technology to be deployed until BLM makes a decision about the right-of-way. Utility Corridor DEIS at ES-2. Enefit's public statements and applications to BLM indicate that it will operate the South Project regardless of whether the right-of-way is granted. Utility Corridor DEIS at ES-2. And Enefit states that it plans

to construct a proven retort technology called Enefit280 on the South Project property as part of its Utah operations. Enefit American Oil, RD&D Development Plan, July 19, 2012 at 10-11.

As you are aware, NEPA obligates reasonable forecasting. In discussing this obligation, courts have found that “the agency need not speculate about all conceivable impacts, but it must evaluate the reasonably foreseeable significant effects of the proposed action.” *Sierra Club v. Marsh*, 976 F.2d 763, 767 (1st Cir. 1992). In this context, reasonable foreseeability means that “the impact is sufficiently likely to occur that a person of ordinary prudence would take it into account in reaching a decision.” *Id.* An environmental effect would be considered “too speculative” for inclusion in the EIS if it cannot be described at the time the EIS is drafted with sufficient specificity to make its consideration useful to a reasonable decision-maker. *Id.* at 768. Nevertheless, “[r]easonable forecasting . . . is . . . implicit in NEPA, and we must reject any attempt by agencies to shirk their responsibilities under NEPA by labeling any and all discussion of future environmental effects as ‘crystal ball inquiry.’” *Scientists’ Inst. for Pub. Info. v. Atomic Energy Comm’n*, 481 F.2d 1079, 1092 (D.C. Cir. 1973).

## **II. BLM Failed to Consider Data Relevant to Enefit’s Oil Shale Operations**

BLM can, using publically available data, forecast air, greenhouse gas, waste, and other environmental impacts based on Enefit’s current use of Enefit280 technology in Estonia and other available data regarding oil shale impacts.

### ***a. The Enefit280 Process***

Enefit has stated that it intends to use a process known as Enefit280 to convert oil shale to synthetic crude oil at its South Project in Utah.<sup>1</sup> This is technology that Enefit touts on its website for its Utah operations as “proven” because the company has operated a retort facility using the Enefit280 technology in Estonia since approximately 2012. Eesti Energia details the results of the Enefit280 operation in its 2016 Q1 interim financial report: “During the quarter, our new Enefit280 oil plant increased its output to 38 thousand tonnes and for the first time contributed more than half of our total shale oil output.”<sup>2</sup> Additionally, there is ample scholarship, reports, and available data focused on oil shale mining impacts both in Estonia and in the United States that BLM should have considered in the draft EIS.

There are extensive studies on the use of Enefit280 technology in Estonia and the oil shale industry in Estonia. Enefit has contracted studies on the Enefit280 technology sufficient to allow them to estimate carbon emissions from the Enefit280 technology. In 2010, Enefit commissioned Jacobs Consultancy to perform a study focused on Enefit’s CO<sub>2</sub> intensity,<sup>3</sup> which evaluated the energy return on investment of the Enefit shale oil industry in Estonia and Enefit’s emissions. In its corporate and social responsibility report of 2014, Eesti Energia states that it has performed or commissioned “studies to investigate the environmental impacts of oil shale mining and processing...in order to compare the benefits of oil shale industry with costs arising from

---

<sup>1</sup> Enefit American Oil, *RD&D Development Plan*, July 19, 2012 at 10-11; *see also*, Hrenko, R., Enefit American Oil: Introduction and Progress Update, University of Utah Unconventional Fuel Conference, Presentation. May 14, 2012. Salt Lake City, Utah. *available at* <http://content.lib.utah.edu/utis/getfile/collection/ir-eua/id/3190/filename/3177.pdf>

<sup>2</sup> Eesti Energia, Q1 2016 Interim Report 1 January 2016 – 31 March 2016, at 4, *available at* [https://www.energia.ee/-/doc/10187/pdf/concern/Interim\\_report\\_2016\\_Q1\\_eng.pdf](https://www.energia.ee/-/doc/10187/pdf/concern/Interim_report_2016_Q1_eng.pdf).

<sup>3</sup> Aarna, I., Lauringson, T. Carbon intensity, water use and EROI of production of upgraded shale oil products using the Enefit280 technology. October 2011. Presentation, Golden, CO. *available at* [http://www.costar-mines.org/oss/31/F-pres-sm-sec/12-4\\_Aarna\\_Indrek.pdf](http://www.costar-mines.org/oss/31/F-pres-sm-sec/12-4_Aarna_Indrek.pdf)

environmental impacts”; and assessments of “the environment impacts of [oil shale] mining including the impact of mining on fauna, water level and groundwater.”<sup>4</sup> Additionally, the air emissions, carbon output, and waste disposal of the Estonian oil shale industry have all been measured and studied by scientists, academics, and regulators.<sup>5</sup>

Enefit has also provided information about the use of the Enefit 280 technology in Utah, including data as detailed as process diagrams, in its investor reports and promotional material.<sup>6</sup> And if Enefit can estimate the number of employees, volume of water required, and tax revenues likely to result from the South Project, as it does in the draft EIS and elsewhere, it can and must generally disclose air, climate, and waste impacts.

The draft EIS’s lack of details on environmental impacts of the South Project, which results largely from Enefit alleging that it has not yet identified exact project parameters, is particularly troubling given that the environmental challenges associated with ex situ mining and the operation of surface retorts in Estonia have been thoroughly studied. Similarly, in the United States, the few limited ex situ operations that did operate produced a range of scholarship that documented a number of environmental challenges associated with these types of developments.

### ***b. Solid and Hazardous Waste Impacts***

A significant environmental impact of mining and processing of oil shale is that it creates substantially more solid waste by volume than what is originally mined.<sup>7</sup> More troubling is that after the shale is retorted, the residual char, or spent shale, is chemically altered for the worse. The spent shale, transformed due to its exposure to increased temperatures, contains a number of soluble inorganics including significant quantities of arsenic and selenium.<sup>8</sup> Compounding matters, spent shale also contains highly carcinogenic polycyclic aromatic hydrocarbons (PAHs).<sup>9,10</sup> In fact, toxic levels of PAHs were found in Green River Basin spent shale that was produced in the early 1980’s.<sup>11</sup>

---

<sup>4</sup> Eesti Energia, Corporate and Social Responsibility Report, 2014, available at [https://www.energia.ee/-/doc/10187/pdf/concern/keskkonnaaruanne\\_2014\\_eng.pdf](https://www.energia.ee/-/doc/10187/pdf/concern/keskkonnaaruanne_2014_eng.pdf).

<sup>5</sup> See, e.g., Gavrilova et al., *Life Cycle Analysis of the Estonian Oil Shale Industry*, Tallin University of Technology, 2005; Brandt, A. *Greenhouse gas emissions from liquid fuels produced from Estonian oil shale*. January 2011. Prepared for European Commission - Joint Research Center, available at [https://circabc.europa.eu/sd/d/9ab55170-dc88-4dcb-b2d6-e7e7ba59d8c3/Brandt\\_Estonian\\_Oil\\_Shale\\_Final.pdf](https://circabc.europa.eu/sd/d/9ab55170-dc88-4dcb-b2d6-e7e7ba59d8c3/Brandt_Estonian_Oil_Shale_Final.pdf)

<sup>6</sup> See, e.g., Enefit. Presentation, available at <http://content.lib.utah.edu/utills/getfile/collection/ir-eua/id/3257/filename/3267.pdf>; Enefit, *Retorting Enefit 280*, <https://www.enefit.com/retorting-enefit280>

<sup>7</sup> The BLM conservatively estimated in the 2007 oil shale PEIS that “Regardless of the retort, spent shale volume would increase by 30%.” BLM. Draft: OSTIS PEIS. 2007d. 4.1.1, P. 4-6.

<sup>8</sup> *Id.*, 4.14.1.2, P. 4-167.

<sup>9</sup> Irha, N.; E. Teinmaa. *Behavior of Three- to Four-Ring PAHs in the Presence of Oil Shale Ash and Aluminosilicate Matter*. 2002. Polycyclic Aromatic Compounds. Volume 22. P. 663 - 671.

<sup>10</sup> Enefit has made claims in various forums and within promotional materials that waste issues will not be as acute with the Enefit 280 process, but the company has not produced any academic studies or other rigorous documentation that substantiates such a prediction.

<sup>11</sup> IARC. Polynuclear Aromatic Compounds, Part 4, *Bitumens, Coal-Tars and Derived Products, Shale-Oils and Soots, Summary of Data Reported and Evaluation, on the Evaluation of Carcinogenic Risks to Humans*, World Health Organization, last updated April 20, 1998. 1985, Volume 35. available at <http://monographs.iarc.fr/ENG/Monographs/vol35/volume35.pdf>.

Even under the best of circumstances, it is not technically evident that one can fully segregate the hazardous char waste stream from the rest of the retorted spent shale material.<sup>12</sup> The inability to separate or manage for these mixed waste streams presents additional challenges. Intrusion and exposure to water concentrates undesirable inorganic elements into quantities that pose critical problems for the overall welfare of an ecosystem.<sup>13</sup> Given the generous volume of wastes produced by a retort facility, the sheer industrial scale of such an operation presents considerable challenges in any endeavor to stabilize and manage such a waste stream. Preventing leaching of inorganic elements in a spent shale waste pile has so far proven to be a practical impossibility.

The abandoned Anvil Points retorting facility near Rifle, Colorado presents a case in point. The experimental retort facility processed shale from 1947 to 1984. During its run, Anvil Points created 61 tons of spent shale.<sup>14</sup> This modest amount of spent shale would be dwarfed by what has been proposed by Enefit. In 2007, BLM estimated that a facility of the size being proposed by Enefit, with the expressed goal of a 50,000 bbl/d, would produce upwards of 23 million tons of spent shale waste a year.<sup>15</sup>

It has been decades since the Anvil Points facility was abandoned, but those 60 tons have been leaching a number of critical inorganic elements into the region's surface water. Foremost in the Anvil Points' leachate is the presence of arsenic - created during the retorting process - that continues to discharge at quantities exceeding Colorado Water Quality Standards.<sup>16</sup> The mere existence of 60 tons of spent shale waste has become a significant environmental and financial liability for the state of Colorado and the federal government. Nearly \$65 million dollars have been allocated to remediate the spent shale waste pile and the surrounding site.

Another challenge is the ignitability of spent shale waste piles. Across the planet, a number of oil shale operations have been plagued by the combustibility of spent shale waste piles.<sup>17</sup> In Colorado, at the aforementioned Anvil Points spent shale pile, a 1995 Department of Energy site report found that, "The shale pile has been self-retorting intermittently since at least 1979, and that heat and vapors from this *in situ* combustion are generated."<sup>18</sup> The Enefit right-of-way draft EIS does not address these impacts.

---

<sup>12</sup> Karhu, A. *Environmental hazard of the waste streams of Estonian oil shale industry*. 2006. Oil Shale. Vol 23. P. 53-93.

<sup>13</sup> Argonne National Laboratory. *Environmental Consequences of, and Control Processes For, Energy Technologies*. 1990. P. 102-115.

<sup>14</sup> Meade, R.B. *Tailings and Mine Waste '02: Proceedings of the Ninth International Conference on Tailings and Mine Waste*, Fort Collins, Colorado, USA, 27-30 January 2002. P. 428.

<sup>15</sup> BLM. Draft: OSTTS PEIS. 4.9.1.1.2 4- p. 119. Notably, the Enefit right-of-way draft EIS fails to contain any such estimate.

<sup>16</sup> Besides critically elevated levels of arsenic, other inorganic elements leaching from the Anvil Points spent shale waste site include: aluminum, arsenic, boron, barium, chromium, cobalt, copper, iron, lithium, magnesium, manganese, molybdenum, sodium, nickel, lead, vanadium, and zinc. See: BLM. Hazardous Materials Management/Abandoned Mine Land Management Applicable or Relevant and Appropriate Requirements. 2007. TR-1703-1/TR-3720-1. P. 23.

<sup>17</sup> A United Nations assessment of oil shale found one spent shale pile in Kumla, Sweden had been on fire since 1947. See, MacDonald M. E., M. J. Chadwick, G. S. Aslanian, B. S. Beloselsky. *The Environmental Management of Low-grade Fuels*. United Nations Environment Programme by the Stockholm Environment Institute. 1996. P. 137.

<sup>18</sup> Department of Energy Naval Petroleum & Oil Shale Reserves. *Site Environmental Report*. 1995. DOE/FE/62316-T1. P.8

Again, the BLM in its PEIS review expressed that there was a significant degree of uncertainty regarding the agency's wherewithal to properly manage and contain spent shale given the number of unknown issues:

Regardless of the disposal option selected, a number of issues need to be addressed, including the structural integrity of emplaced spent shale, an increase in volume (and decrease in density) over raw shale during the retorting process (this has become known as "the popcorn effect"), and the character of leachates from spent shale. Limited research has been conducted on each of these issues (*emphasis added*).<sup>19</sup>

And:

Field data evaluating the leachate character of spent shale have been collected by the EPA and others. Although the data are limited, there appears to be a clear indication that subjecting oil shale to retorting conditions can result in the mobilization of various ionic constituents contained in the mineral portion of the oil shale.<sup>20</sup>

The EPA, in 2007, also expressed concern about the lack of empirical data necessary to ensure that spent shale waste could be properly managed:

Waste issues are important, particularly for surface retort methods. EPA is currently coordinating with a number of entities on sampling and analytical protocols regarding wastes derived from retort methods. This is currently a high priority for EPA.<sup>21</sup>

Possibly most telling are the measures that the European Union (EU) have taken to further tighten the regulatory controls that govern the disposition of spent shale as a hazardous material in Estonia. In 2000, facing the inclusion of Estonia as a new member of the EU, the EU adopted increasingly more stringent requirements for the management of spent shale waste.<sup>22</sup> The EU was motivated to act given that Estonia had generated over 110 million tons of spent shale waste (generated from aboveground retorting of oil shale). In 2003, after further analysis revealed that the spent shale waste piles created by the Estonian oil shale industry were exceedingly toxic, the EU issued specific guidance to further regulate the administration of spent shale wastes created by retorting.<sup>23</sup>

---

<sup>19</sup> BLM. Draft: OSTs PEIS. A-4, P. A-49.

<sup>20</sup> *Id.*

<sup>21</sup> Hogle, D. 2007 Oil Shale Environmental Issues and Needs Workshop October 18, 2007, Colorado School of Mines, Golden, Colorado. U.S. Department of Energy-Office of Fossil Energy-National Energy Technology Laboratory-Strategic Center for Natural Gas & Oil. 2008. P. 12.

<sup>22</sup> 2000/532/EC: Commission Decision of 3 May 2000 replacing Decision 94/3/EC establishing a list of wastes pursuant to Article 1(a) of Council Directive 75/442/EEC on waste and Council Decision 94/904/EC establishing a list of hazardous waste pursuant to Article 1(4) of Council Directive 91/689/EEC on hazardous waste (notified under document number C(2000) 1147) OJ L 226 , 06/09/2000 P. 0003–0024.

<sup>23</sup> Council Decision 2003/33/EC Establishing criteria and procedures for the acceptance of waste at landfills pursuant to Article 16 and Annex II to Directive 1999/31/EC , OJ L 11/27 , 16.1.2003.

One anecdotal byproduct of increased regulation by the EU was a notable escalation in the number of scientific studies that investigated the hazardous nature of spent shale. Many of these additional analyses originated from Estonia, where past practices by Estonia's oil shale industry had created ample opportunity to scrutinize the impacts of spent shale waste.

Despite this wealth of information, the Enefit right-of-way draft EIS merely acknowledges that there may be some kinds of impacts; declares that such impacts can't be predicted; or declares that whatever the impacts are, they are irrelevant.<sup>24</sup> This lack of analysis violates NEPA.

### ***c. Climate Change and Air Quality Impacts***

Besides the spent shale waste issue that has plagued Estonia, the processing of oil shale into electricity and petroleum products has had profound environmental implications in the context of climate change. A number of papers have established that CO<sub>2</sub> emissions from oil shale processing are possibly the dirtiest feedstock to be found on the planet.<sup>25</sup> Even Enefit's promotional materials regarding emission factors – which are based on a number of optimistic or at least unchallenged assumptions – claim that the CO<sub>2</sub> emissions of the Enefit 280 process will still be dirtier than current conventional fuel development or even tar sands.<sup>26</sup>

Our organizations understand that it is likely that the Utah operation will, due to geology and design, not be *exactly* the same as Enefit's operations in Estonia. However, Enefit's Estonian operations are clearly models for what Enefit will construct in Utah. Therefore, available studies, reports and other available information provide a useful metric for the public and decision-makers to understand the likely nature and scale of impacts to air quality and climate, something now completely omitted from the draft EIS, in violation of NEPA.

## **III. Conclusion.**

The South Project and Enefit's plans to expand its mining activities onto federal and state land are both connected actions and cumulative impacts of the right-of-way decision. As part of its NEPA obligations, BLM should comply with its reasonable forecasting obligations using the existing data on of the Enefit280 technology and the impacts of oil shale in Estonia and in the United States in its analysis in the draft EIS. BLM's failure to do so violates NEPA regulations

---

<sup>24</sup> See Enefit Right-of-Way Draft EIS at 4-68 (stating only vaguely that "Spent shale piles and mine tailings ... might be sources of contamination for salts, metals, and hydrocarbons for both surface and groundwater"); *id.* at 4-70, 4-72, and 4-94 (making similar statements). See also *id.* at 4-138 (declining to disclose any information about the public health or other impacts of spent shale, alleging that such data is "unknown, and cannot be obtained, due to the fact that design and engineering of the South Project will change based on whether or not the BLM allows the Applicant to build one or more of the proposed utilities. BLM believes this unknown information is not essential to a reasoned choice between alternatives").

<sup>25</sup> Brandt, A. *Greenhouse gas emissions from liquid fuels produced from Estonian oil shale*. January 2011. Prepared for European Commission - Joint Research Center, available at [https://circabc.europa.eu/sd/d/9ab55170-dc88-4dcb-b2d6-e7e7ba59d8c3/Brandt\\_Estonian\\_Oil\\_Shale\\_Final.pdf](https://circabc.europa.eu/sd/d/9ab55170-dc88-4dcb-b2d6-e7e7ba59d8c3/Brandt_Estonian_Oil_Shale_Final.pdf); Mui, S., Tonachel, L., McEnaney, B., Shope, E. *GHG Emission Factors for High Carbon Intensity Crude Oils*. 2010. Natural Resources Defense Council, available at [https://www.nrdc.org/sites/default/files/ene\\_10070101a.pdf](https://www.nrdc.org/sites/default/files/ene_10070101a.pdf).

<sup>26</sup> Aarna, I., Lauringson, T. *Carbon intensity, water use and EROI of production of upgraded shale oil products using the Enefit280 technology*. October 2011. Presentation, Golden, CO, available at [http://www.costar-mines.org/oss/31/F-pres-sm-sec/12-4\\_Aarna\\_Indrek.pdf](http://www.costar-mines.org/oss/31/F-pres-sm-sec/12-4_Aarna_Indrek.pdf).

and deprives the public of the opportunity to understand Enefit's impact on public health and the environment.

Developing the oil shale industry in the United States will have significant impacts on our land, species, water, and climate. Yet, if the status quo continues, Enefit will begin operations of the first commercial-scale oil shale project in the United States under a flawed NEPA review that fails to consider available and relevant information about the environmental and public health impacts of oil shale mining and processing. We urge EPA to exert its influence as a regulatory agency and coordinating agency to improve the ongoing NEPA process.

We attach a list of referenced studies, any of which we are happy to provide at your request.

Sincerely,

Edward B. Zukoski,  
Staff Attorney  
Earthjustice, Rocky Mountain Office  
633 17th Street, Suite 1600  
Denver, CO 80202  
(303) 996.9622

Bobby McEnaney  
Senior Deputy Director  
Western Renewable Energy Project  
Natural Resource Defense Council  
1152 15th Street NW Suite 300.  
Washington, D.C. 20005  
(202) 289.6868.

Anne Mariah Tapp  
Energy Program Director  
Grand Canyon Trust  
2601 N. Fort Valley Rd  
Flagstaff, AZ 86001  
(512) 565.9906

Rob DuBuc  
Staff Attorney  
Western Resource Advocates  
150 South 600 East, Ste 2A  
Salt Lake City, Utah 84102  
(801) 487.9911

Steve Bloch  
Legal Director  
Southern Utah Wilderness Alliance  
425 East 100 South  
Salt Lake City, UT 84111  
(801) 428.3981

## LIST OF REFERENCES

Aarna, I., Lauringson, T. Carbon intensity, water use and EROI of production of upgraded shale oil products using the Enefit280 technology. October 2011. Presentation, Golden, CO.  
[http://www.costar-mines.org/oss/31/F-pres-sm-sec/12-4\\_Aarna\\_Indrek.pdf](http://www.costar-mines.org/oss/31/F-pres-sm-sec/12-4_Aarna_Indrek.pdf)

Argonne National Laboratory. Environmental Consequences of, and Control Processes For, Energy Technologies. 1990. P. 102-115.

Arro, H., A. Prikk, T. Pihu, I. Opik. Utilization of semi-coke of Estonian shale oil industry. 2002. Oil Shale. Vol. 19. P. 117-125.

Aunela-Tapola, L., Frandsen, F., Häsänen, E. Trace metal emissions from the Estonian oil shale fired power plant. August 1998. Fuel Processing Technology. Volume 57, Issue 1. P. 1-24

Barton, T., B. Braunberger. Mercury in Oil Shale, Problems and Solutions. Colorado School of Mines, 28th Oil Shale Symposium, October 13-17, 2008.

Bell, P. R. F., Krol, A. A., Greenfield, P. F. Factors controlling the leaching of major and minor constituents from processed Rundle oil shale. 1986. Water Res. Vol. P. 741-750.

Bispo, A., Jourdain, M. J., Jauzein, M. Toxicity and genotoxicity of industrial soils polluted by polycyclic aromatic hydrocarbons (PAHs). 1999. Organ. Geochem. Vol. 30. P. 947-952.

Bitjukova, L., Sorlie, J.-E., Saether, Ola M., Petersell, V. Estimation of organic pollutants in groundwater near oil shale waste mounds in Kohtla-Järve (North-Eastern Estonia). COGEOENVIRONMENT, 2003. International Workshop. Geosciences for urban development and environmental planning. September 13-18, 2003. Vilnius. Extended abstracts. P. 15-17.

Brandt, A. Greenhouse gas emissions from liquid fuels produced from Estonian oil shale. January 2011. Prepared for European Commission - Joint Research Center.

Brendow, K. Global oil shale issues and perspectives. 2003. Oil Shale. Vol. 20. P. 81-92.

Bureau of Land Management. Hazardous Materials Management/Abandoned Mine Land Management Applicable or Relevant and Appropriate Requirements. 2007. TR-1703-1/TR-3720-1. P. 23.

Connell, L.D., P.R. Bell. Description of a Field Experiment To Measure Leachate Formation From Oil Shale Wastes. Waste Management & Research. 1992. Vol. 10. P. 517-533.

Department of Energy Naval Petroleum & Oil Shale Reserves. Site Environmental Report. 1995. DOE/FE/62316-T1. P. 8.

Fox, J.P. Leaching of oil shale solid wastes: A critical review. Center for Environmental Sciences. 1983.



Gavrilova et al., *Life Cycle Analysis of the Estonian Oil Shale Industry*, Tallin Univeristy of Technology, 2005

Gavrilova, O., Vilu, R., Vallner, L. A life cycle environmental impact assessment of oil shale produced and consumed in Estonia. December 2010. *Resources, Conservation and Recycling*. Volume 55, Issue 2. P. 232–245.

Grosse, D.W., L. McGowan. *Tar Sands Leachate Study*. 1984. EPA 600/2-84-113, U.S. Environmental Protection Agency, Industrial Environmental Research Laboratory, Cincinnati, Ohio, Feb.

Habicht J, Kaja Orupõld. *Retorted Oil Shale In Estonia: An Environmental Audit*. Colorado School of Mines, 28th Oil Shale Symposium, October 13-17, 2008.

HARZIA, H., K. ORUPÕLD, J. HABICHT, T. TENNO. Leaching behavior of oil shale semicoke: sulphur species. 2007. *Oil Shale*. Vol. 24. P. 583-589.

Hogle, D. 2007 Oil Shale Environmental Issues and Needs Workshop October 18, 2007 Colorado School of Mines, Golden, Colorado. U.S. Department of Energy-Office of Fossil Energy-National Energy Technology Laboratory-Strategic Center for Natural Gas & Oil. 2008. P. 12.

IARC. *Polynuclear Aromatic Compounds, Part 4, Bitumens, Coal-Tars and Derived Products, Shale-Oils and Soots, Summary of Data Reported and Evaluation, on the Evaluation of Carcinogenic Risks to Humans*, World Health Organization, last updated April 20, 1998. 1985, Volume 35.

Irha, N, E. Teinmaa. Behavior of Three- to Four-Ring PAHs in the Presence of Oil Shale Ash and Aluminosilicate Matter. 2002. *Polycyclic Aromatic Compounds*. Volume 22. P. 663 - 671.

Jabe, J. O., S. D. Probert, M. A. Tahat. Predicted environmental and social impacts of the proposed oil shale integrated tri-generation system (OSITGS). 1999. *Oil Shale*. Vol. 16. P. 2-29.

Kahru, A., M. Kurvet, I. Kurvet. Study of the toxicological impact of different components of ash-heap water (sulphur rich phenolic leachate) using luminescent bacteria as test organisms. 1997. *Oil Shale*. Vol. 14. P. 469-475.

Kahru, A., Maloverjan, A., Sillak, H., Põllumaa, L. Toxicity and fate of phenolic pollutants in the contaminated soils associated with the oil-shale industry. 2002. *Environ. Sci. Pollut. Res*. Vol. 1. P. 27–33.

Kahru, A., Põllumaa, L., Reiman, R., Rätsep, A. *Microbiotests for the evaluation of the pollution from the oil shale industry*. New York, USA: Kluwer Academic/Plenum Publishers, 2000. P. 357–365.

- Kahru, A., Põllumaa, L., Reiman, R., Rätsep, A. Predicting the toxicity of oil shale industry wastewater by its phenolic composition. 1999. *ATLA*. Vol. 27. P. 359–366.
- Kahru, A., Põllumaa, L. Environmental hazard of the waste streams of Estonian oil shale industry: an ecotoxicological review. 2006. *Oil Shale*. Vol. 23. P. 53–93.
- Laja, M., N. Irha, , G. Urb, , J. Reinik, U. Kirso. Characterization of waste extracts of solid waste generated by oil shale combustion. 2005. *Proc. Estonian Acad. Sci. Chem.* Vol. 54. P. 134–141.
- Laja, M., N. Irha, , G. Urb, , J. Reinik, U. Kirso. Leaching behaviour of ash fractions from oil shale combustion by fluidized bed and pulverized firing processes. 2005. *Oil Shale*. Vol. 22. P. 453–465.
- MacDonald, M. E., M. J. Chadwick, G. S. Aslanian, B. S. Beloselsky. *The Environmental Management of Low-grade Fuels*. United Nations Environment Programme by the Stockholm Environment Institute. 1996. P. 125.
- Maremäe, E., V. Ahelik. Leaching of metals from Baisun oil shale and its ashes under the effect of sulphuric acid at low content of liquid phase. 1992. *Oil Shale*. Vol. 9. P. 138-144.
- Maremäe, E., I. Johannes, V. Ahelik. The percolative leaching of alum shale's sulphated ashes. *Oil Shale*. 1991. Vol. 8. P. 138-147.
- McWhorter, D., D. Durnford. **LEACHING AND HYDRAULIC PROPERTIES OF RETORTED OIL SHALE INCLUDING EFFECTS FROM CODISPOSAL OF WASTEWATER**. U.S. Environmental Protection Agency, Washington, D.C., EPA/600/7-86/035.
- Meade, R.B. *Tailings and Mine Waste '02: Proceedings of the Ninth International Conference on Tailings and Mine Waste*, Fort Collins, Colorado, USA, 27-30 January 2002. P. 428.
- Meyer, J.S., D.A. Sanchez, J.A. Brookman, D.B. McWhorter, H.L. Bergman. **CHEMISTRY AND AQUATIC TOXICITY OF RAW OIL SHALE LEACHATES FROM PICEANCE BASIN, COLORADO**. 1985. *Environmental Toxicology and Chemistry*. Vol. 4. P. 559–572.
- Meyer, J. Contribution of Major Ions to Toxicity of Oil Shale Leachates to Aquatic Organisms. Colorado School of Mines, 28th Oil Shale Symposium, October 13-17, 2008.
- Mölder, L., Elenurm, A., Tamvelius, H. Sulphur compounds in a hydraulic ash-disposal system. 1995. *Proc. Estonian Acad. Sci. Chem.* Vol.44. P. 207–211.
- Motlep, R. Mineral Composition of Estonian Oil Shale Semi-coke Sediments. 2007. *Oil Shale*. Vol. 24. P. 405–422.

- Mui, S., Tonachel, L., McEnaney, B., Shope, E. GHG Emission Factors for High Carbon Intensity Crude Oils. 2010. Natural Resources Defense Council.  
[https://www.nrdc.org/sites/default/files/ene\\_10070101a.pdf](https://www.nrdc.org/sites/default/files/ene_10070101a.pdf)
- Munter, R., Trapido, M., Veressinina, J., Preis, S., Pikkov, L. Detoxification of phenolic and carcinogenic compounds in the leachate from Kohtla-Järve dump. Proceedings of Estonian Conference on Ecology. Tartu, Estonia: Greif, 1994. P. 199–203.
- Nataly, I, E. Teinemaa. Behavior of Three- to Four-Ring PAHs in the Presence of Oil Shale Ash and Aluminosilicate Matter. Polycyclic Aromatic Compounds. 2002. Volume 22. P. 663 – 671.
- Naumov, B. Pollution of hydrosphere by leaching of dictyonema shale in waste dumps of the Maardu openpits. 1991. Oil Shale. Vol. 8. P. 266-274.
- Orupõld, K., Ohlsson, A., Henrysson, T. Batch trials to simulate biological treatment in lagoons of leachate from oil-shale ash heaps. 1997. Oil Shale. Vol. 14. P. 476–487.
- Orupõld, K., Tenno, T., Henrysson, T. Biological lagooning of phenols-containing oil shale ash heaps leachate. 2000. Water Res. Vol. 34. P. 4389–4396.
- Orupõld, K.; Habicht, J.; Tenno, T. Leaching behaviour of oil shale semicoke: compliance with the waste acceptance criteria for landfills. 2008. Oil Shale. Vol. 25. P. 267 - 275.
- Otsa, E., Tang, H. The determining of environmental hazard of semi-coke. The report of the Estonian Environmental Research Centre.-Tallinn, 2003.
- Parakhonsky E. Theoretical and technological principles of preventing spontaneous ignition of oil shale mines waste heaps. 1992. Oil Shale. Vol. 9. P. 346-356.
- Pereira, W.E., C. Rostad. Chemical Characterization of Organic Compounds in Leachates from Surface-Retorted Oil Shale Generated by Gas Combustion and Indirect Pyrolysis. 1983. International Journal of Environmental Analytical Chemistry, Volume 15, P. 73 – 88.
- Põllumaa, L., Maloveryan, A., Trapido, M., Sillak, H., Kahru, A. Study of the Environmental Hazard Caused by the Oil-Shale Industry Solid Waste. 2001. ATLA. Vol. 29. P. 259–267.
- Ratsep, A., E. Rull, V. Liblik. Impact of oil shale mine water discharges on phytoplankton community of Purtse catchment rivers. 2002. Oil Shale. Vol. 19. P. 297-306.
- Roots, O. Polychlorinated biphenyls (PCB), polychlorinated dibenzo-p-dioxins (PCDD) and dibenzofurans (PCDF) in oil shale and fly ash from oil shale-fired power plant in Estonia. 2004. Oil Shale. Vol. 21. P. 333–339.

- Soone, J., S. Doilov. Sustainable utilization of oil shale resources and comparison of contemporary technologies used for oil shale processing. 2003. *Oil Shale* Vol. 20, No. 3 Special Symposium on Oil Shale, 18-21 November 2002. P. 311-323.
- Sorlie, J.E. Risk based environmental site assessment of waste from the oil shale industry. Integration-4-Water Workshops Event, 7th Framework Programme of the European Union, Warsaw, Poland. 2-4 June 2006.
- Stollenwerk, K.G., D.D. Rundells. Composition of leachate from surface-retorted and unretorted Colorado oil shale. 1981. *Environment Science & Technology*. Vol. 15. P. 1340-1346.
- Tamm, K., et al. LEACHING THERMODYNAMICS AND KINETICS OF OIL SHALE WASTE KEY COMPONENTS. 2016. *Oil Shale*. Vol. 33, Issue 1. P. 80–99.
- Trapido, M.; R. Munter; Y. Veressinina; N. Kulik. Oil Shale Semicoke Leachate Treatment using Ozonation and the Fenton Oxidation. 2006. *Environmental Technology*. Volume 27. P. 307 – 315.
- Trikkel, A., R. Kuusik, N. Maljukova. Distribution of organic and inorganic ingredients in Estonian oil shale semicoke. 2004. *Oil Shale*. Vol. 21. P. 227-235.
- Tuvikene, A., Huuskonen, S., Koponen, K., Ritola, O., Mauer, Ü., Lindström-Seppä, P. Oil shale processing as a source of aquatic pollution: monitoring of the biologic effects in caged and feral freshwater fish. 1999. *Environ. Health Perspect.* Vol. 107. P. 745–752.