

Live Storage Capacity of All Colorado River Basin Reservoirs by Percentage and by Decade

| Water Year | Capacity % | Water Year | Capacity % | Water Year | Capacity % | Water Year | Capacity % |
|----------------|-----------------------------------------------------------------------------------------------------|------------|--------------|------------|--------------|------------|--------------|
| 1980 | 90.3 | 1990 | 72.76 | 2000 | 85.1 | 2010 | 55.46 |
| 1981 | 82.07 | 1991 | 70.75 | 2001 | 77.87 | 2011 | 64.86 |
| 1982 | 89.49 | 1992 | 69.21 | 2002 | 63.54 | 2012 | 57.05 |
| 1983 | 97.74 | 1993 | 81.2 | 2003 | 57.13 | 2013 | 50.21 |
| 1984 | 95.03 | 1994 | 75.48 | 2004 | 50.03 | 2014 | 50.37 |
| 1985 | 91.98 | 1995 | 86.36 | 2005 | 58.59 | 2015 | 50.83 |
| 1986 | 92.14 | 1996 | 84.83 | 2006 | 56.21 | 2016 | 50.62 |
| 1987 | 90.98 | 1997 | 92.53 | 2007 | 53.88 | 2017 | 55.2 |
| 1988 | 87 | 1998 | 93.65 | 2008 | 57.12 | 2018 | 46.97 |
| 1989 | 80.38 | 1999 | 93.65 | 2009 | 57.38 | 2019 | |
| Average | 89.65 | | 82.04 | | 61.69 | | 53.51 |
| Note: | Impacts to hydropower production may occur at capacities near or below 35 % | | | | | | |
| Note: | Impacts of low capacity reservoirs include degradation to water quality | | | | | | |
| Note: | 1980 is the year that Lake Powell filled for the first time; filling of the reservoir began in 1963 | | | | | | |